

Research Article

Preliminary inventory of metazoan parasites of the Lessepsian bluespotted cornetfish *Fistularia commersonii* in the Gulf of Gabès, Tunisia

Wiem BOUSSELLAA ^{*1, 2} , Hela DERBEL ¹  & Lassad NEIFAR ¹ 

¹ Sfax University. Faculty of Science of Sfax. Department of Life Sciences, Sfax, Tunisia

² Marine Turtle Research Group, Department of Marine Animal Conservation and Public Engagement, Stazione Zoologica Anton Dohrn, Via Nuova Macello 16, 80055 Portici, Italy

*Correspondence: wiem.bousellaa@szn.it

Received: 02/05/2023; Accepted: 29/10/2023; Published: 23/11/2023

Abstract: Biological invasions in the world are widely acknowledged as a serious threat to biodiversity, but the role of parasites in marine ecosystem processes is still poorly understood. Among these invasions we are interested in the present work in the parasitofauna of bluespotted cornetfish *Fistularia commersonii* introduced species through the Suez Canal from the Red Sea to the Mediterranean Sea and so-called Lessepsian migration. 43 specimens of *F. commersonii* were sampled from the Gulf of Gabès. Parasite prevalence and intensity were investigated. Moreover, we reviewed the literature to identify native and invasive parasite species recorded. Eleven parasite species infecting *Fistularia commersonii* from the Gulf of Gabès were found.

Our results suggest the co-introduction of five parasite species one Acanthocephala *Sclerocollum* sp., four Digenea *Allolepidapedon petimba*, *Neoallopepidapedon hawaiiense*, *Ectenurus virgulus*; Stephanostomum sp. that are assumed to be originate from the Red Sea. In addition, six parasites have been acquired from the Mediterranean Sea two Cestoda Phyllobothriidae and *Nybelinia Africana*, one Digenea *Lecithochirium* sp., one Hirudinea *Trachelobdella lubrica*, one Isopoda *Gnathia* sp. and one Nematoda *Hysterothylacium* sp. These results show the importance of using parasite community studies, such as the one described here, to understand the role of parasites in Lessepsian migration.

Keywords: Metazoan parasites; Lessepsian migration; *Fistularia commersonii*; Gulf of Gabès; Tunisia; Mediterranean.

1. Introduction

The influx of Red Sea species into the Mediterranean Sea via the Suez Canal has been well documented (Golani, 1998). Among all taxa, fish have been the most studied, due to their commercial importance and their relatively clear taxonomy.

The bluespotted cornetfish, *F. commersonii* is an Indo-Pacific and Red Sea species (Fritzsche, 1976). This species has spread rapidly throughout the Mediterranean to be one of the most successful invaders in the Mediterranean Sea (Streftaris & Zenetos, 2006). Firstly, it was recorded in the Eastern Mediterranean coasts (Golani,

2000) and in a short time, it spread until reaching the Western Basin of the Mediterranean Sea in the Gulf of Sicily (Pipitone et al., 2004), Tunisian coast (Ben Souissi et al., 2004), Sardinia (Pais et al., 2007), and in 2007, it reached the sea of Alboran (Sanchez-Tocino et al., 2007), which is the furthest point than a migrant lessepsien has never been recorded since its introduction (CIESM, 2009). Genetic studies have shown that this successful invasion was the result of a single introduction episode that resulted in a serious bottleneck of the Mediterranean population (Golani et al., 2007).

Many researchers were interested in documenting the presence of this Lessepsian sprinter fish along the Mediterranean coasts. Other investigations focused on the population structure of the species (Bariche, & Kajajian, 2012), biological data (Edelist, 2014; Castriota et al., 2014; Mouine-Oueslatia et al., 2017), diet (Bariche, 2007; Bariche et al., 2009; Kalogirou et al., 2007), reproduction (Bariche et al., 2013), length-weight relationships (Taskavak & Bilecenoglu, 2001; Erguden et al., 2009) and maximum length (Torcu Koc et al., 2019). At present, few data have been published on the parasites of *F. commersonii*. In the Mediterranean Sea, two parasitological study was published by Pais et al. (2007) and Merella et al. (2016), adding new information on the parasite fauna of this migrant fish. However, no studies have been done in the Tunisian coasts. Here we investigated metazoan parasite infections in the Lessepsian fish *F. commersonii* Rüppel, 1838, along the Tunisian coast. By sampling this invasive fish along the Gulf of Gabès, and by conducting an additional parasitological literature survey, we aimed to provide baseline information on the parasite fauna of *F. commersonii* from Tunisian coast.

2. Material and Methods

Between 2014 and 2016, 43 specimens (28 females and 15 males) of *Fistularia commersonii* were collected from different fishing area in the Gulf of Gabès; Tunisia (Figure1).




Figure 1. Sampling Location of *F. commersonii* in the Gulf of Gabès, Tunisia (Central Mediterranean Sea).

Location names, coordinates, sampling dates and sampling effort per species are mentioned in the Table1.

Fish skin, fins, nasal pits, eyes and buccal cavities were thoroughly examined under a stereomicroscope with incident light for the presence of ectoparasites. Gill arches were separated by incision, removed, rinsed and examined individually. Internal organs (stomach, pyloric caeca, intestines, rectum, heart, liver, spleen, gall bladder and gonads) were separated and individually examined for the presence of endoparasites. Parasites were identified to the lowest taxonomic level possible using Gibson et al. (2002) for Digenea, the keys of Kabata (2003) for Copepoda, Berland (1961) and Petter & Maillard (1988) for Nematoda and Euzet (1994) for Cestoda. We searched literature databases for published records of additional parasite species from the Mediterranean Sea and the Red Sea. Parameters of infection such as prevalence (P) and mean intensity (MI)

were determined following Bush et al. (1997).

3. Results and discussions

Based on samples from the Tunisian coasts, eleven parasite species infecting *Fistularia commersonii* were found. (Table2). Our results suggest that the Lessepsian migrant has co-introduced five parasite species to the Tunisian coasts one *Acanthocephala Sclerocollum* sp., four Digenea *Allolepidapedon petimba*, *Neoallopeidapedon hawaiiense*, *Ectenurus virgulus*; *Stephanostomum* sp. and four elsewhere in the Mediterranean Sea (Merella et al., 2016), that are assumed to be originate from the Red Sea. In fact, the two exotic Digenea *A. petimba* and *N. hawaiiense* are the two most prevalent parasites in the Gulf of Gabès (100% and 70%, successively) and the non-native Digenea *A. petimba* has the highest intensity (IM = 62) in the Gulf of Gabès. In spite of their heteroxenous life cycle, these parasites seem to be co-introduced in the Mediterranean Sea. This successful introduction can be explained that parasites had found suitable hosts and environmental conditions to complete the life cycle in the new habitat or their intermediate hosts are already co-introduced in the Mediterranean Sea and their life cycle has been achieved (Boussellaa et al., 2017). However, the fact that all the specimens of these parasites were retrieved at the adult stage could also suggest a residual infection that took place before of the host migration. In addition, we found that the invasive fish has acquired six parasites in the Tunisian coasts two Cestoda Phyllobothriidae and *Nybelinia Africana*, one Digenea *Lecithochirium* sp., one Hirudinea *Trachelobdella lubrica*, one Isopoda *Gnathia* sp., one Nematoda *Hysterothylacium* sp. that are assumed native in the Mediterranean Sea. Introduced fish are naive hosts to native

parasites (Colautti et al. 2004). Therefore, native parasites could represent an important source of selection against migrants (MacColl & Chapman, 2010). The high infection parameters of *F. commersonii* parasites found in the Mediterranean Sea, suggesting a potential competitive advantage and confirm that this species is today one of the most successful invaders of the Mediterranean Sea.

4. Conclusion

According to our research, the Lessepsian migrant may have an impact on native fish hosts by changing the population dynamics of local parasite species through parasite co-introduction, and parasite acquisition, leading to higher infection levels in invaded areas. The success of *F. commersonii* which is the only Lessepsian fish that shows a severe bottleneck in the Mediterranean Sea which offer in the new habitat a rich parasite assemblage that which could potentially infect it even more than in its native area.

Table 1. Information on the sampling design of *Fistularia commersonii* collected for this study from the Gulf of Gabès. Given are the location name, geographic coordinates, sampling dates and sample sizes (per sex).

Location number	Location name	Geographic coordinates	Sampling dates	<i>Fistularia commersonii</i>	
				Females (Size min-max) (cm)	Males (Size min-max) (cm)
1	Kerkennah	34° 39' 29" N 11° 04' 07" E	17/09/2014 16/10/2014 15/02/2016	6 (79.5-87)	2 (74.5-78.3)
2	Skhira	34° 17' 57" N 10° 04' 11" E	13/10/2014	3 (78.5-85.6)	1 (86.8)
3	Zarat	33° 39' 59" N 10° 20' 59" E	05/02/2014 01/12/2014 25/07/2015	8 (82-93.5)	4 (67.5-99)
4	Zarzis	33° 30' 14" N 11° 06' 43" E	25/07/2015	6 (71.7-89.2)	2 (72.4-79.8)
5	El-Keteef	35° 40' 7" N 10° 53' 32" E	17/06/2014 08/11/2014	5 (78.6-90.5)	6 (73.5-85)
Total				28	15

Table 2. Parasite species found in this study and recorded in the literature, with information on the life cycle and the invasive/native status in the Mediterranean Sea and occurrence in *Fistularia commersonii* from the Mediterranean Sea. If quantitative data were available, prevalence and mean intensity ($\pm$ SE) are given.

Parasite taxa	Parasite species	Infection site	Status Med. Sea	<i>Fistularia commersonii</i>		Référence
				P (%)	MI	
Acanthocephala	<i>Sclerocollum</i> sp.	Intestine	invasive	9.3	1.25($\pm$ 0.06)	This study
	<i>Breizacanthus ligur</i>	Intestine	native	17.6	1.5	Merella et al. (2016)
Cestoda	Phyllobothriidae	Intestine	native	41.86	69.11($\pm$ 14.85)	This study
	<i>Nybelinia africana</i>	Stomach	native	30.23	25.38 ($\pm$ 3.58)	This study
	<i>Pseudogrillotia</i> sp.	Intestine	native	23.5	1.9	Merella et al. (2016)
	<i>Dasyrhynchus basipunctatus</i>	Intestine	invasive	-	-	Beveridge et al. (2014)
	<i>Caligus fistulariae</i>	Mouth cavity	invasive	-	-	Merella et al. (2016)
Copepoda	<i>Caligus elongatus</i>	Mouth/gills	native	5.9	1	Merella et al. (2016)
	<i>Caligus minimus</i>	Mouth	native	2.9	1	Merella et al. (2016)
	<i>Caligus flexispina</i>	Mouth	invasive	-	-	Rigby et al. (1999)
	<i>Allolepidapedon petimba</i>	Stomach/Intestine	invasive	100%	61.72($\pm$ 9.81)	This study
Digenea	<i>Neoallolepidapedon hawaiiense</i>	Stomach	invasive	69.76	3.46 ($\pm$ 0.48)	This study
	<i>Ectenurus virgulus</i>	Stomach	invasive	18.60	1($\pm$ 0.06)	This study
	<i>Lecithochirium</i> sp.	Stomach	Native	23.25	1.4 ($\pm$ 0.1)	This study
	<i>Stephanostomum</i> sp.	Intestine	invasive	16.28	1.14($\pm$ 0.07)	This study
	<i>Allolepidapedon fistulariae</i>	Stomach/Intestine	invasive	66.7	7.5	Merella et al. (2016)
	Bucephalidae	Intestine	native	2.9	3	Merella et al. (2016)
	Hemiuridae	Stomach	native	2.9	1	Merella et al. (2016)
	Trematoda (juvenile)	Stomach	native	2.9	1	Merella et al. (2016)
	Trematoda (metacercaria)	Stomach	native	8.8	1.3	Merella et al. (2016)
	<i>Lecithochirium microstomum</i>	Stomach	native	-	-	Rigby et al. (1999)
Hirudinea	<i>Trachelobdella lubrica</i>	Mouth/gills	native	20.93	1 ($\pm$ 0.06)	This study
Isopoda	<i>Gnathia</i> sp.	Mouth/gills	native	13.95	3.16($\pm$ 0.23)	This study
Nematoda	<i>Hysterothylacium</i> sp.	Body cavity/digestive tract	native	69.76	8.73($\pm$ 2)	This study
	<i>Hysterothylacium aduncum</i>	Body cavity/digestive tract	native	8.8	2.3	Merella et al. (2016)
	<i>Hysterothylacium fabri</i>	Body cavity/digestive tract	native	41.2	5.1	Merella et al. (2016)
	<i>Spinitectus</i> sp.	Intestine	native	8.8	1.3	Merella et al. (2016)
	Cystidicolidae	Stomach	native	2.9	1	Merella et al. (2016)
	Philometridae	Stomach	native	2.9	1	Merella et al. (2016)




Figure 2. Parasites collected from *Fistularia commersonii* from the Gulf of Gabès

Acknowledgments

We thank Dr. Maroof Khalaf and Dr. Tariq Al-Najjar from the Faculty of Marine Sciences, Department of Aqaba Marine Biology and all fishermen of Gulf of Gabès for their help during literature investigation and the fish sampling mission. We thank Pr. Lassad Neifar and Dr. Hela Derbel for their useful comments on parasite identification.

References

1. Bariche, M. (2007). Diet of the Lessepsian *Fistularia commersonii* (Teleostei, Fistulariidae) off the coast of Lebanon: preliminary results. *Rapport du 38^e Congrès de la CIESM*, 38, 425.
2. Bariche, M., Alwan, N., El-Assi, H. & Zurayk, R. (2009). Diet composition of the Lessepsian bluespotted cornetfish *Fistularia commersonii* in the eastern Mediterranean. *Journal of Applied Ichthyology*, 25(4), 460-465. <https://doi.org/10.1111/j.1439-0426.2008.01202.x>
3. Bariche, M. & Kajajian, A. (2012). Population structure of the bluespotted cornetfish *Fistularia commersonii* (Osteichthyes: Fistulariidae) in the eastern Mediterranean. *Journal of Biological Research-Thessaloniki*, 17, 74-80.
4. Bariche, M., Kazanjian, G. & Azzurro, E. (2014). A lag of 25 years: evidence from an old capture of *Fistularia commersonii* Rüppell, 1838 from Lebanon (Mediterranean Sea). *Journal of Applied Ichthyology*, 30 (3): 535-536. <https://doi.org/10.1111/jai.12394>
5. Ben Souissi, J., Zaouali, J., Bradai, M.N. & Quignard, J.P. (2004). Lessepsian migrant fishes off the coasts of Tunisia. First record of *Fistularia commersonii* (Osteichthyes, Fistularidae) and *Parexocoetus mento* (Osteichthyes, Exocoetidae). *Vie et Milieu*, 54 (4), 247-248
6. Beveridge, I., Bray, R.A., Cribb, T.H. & Justine J.L. (2014). Diversity of trypanorhynch metacestodes in teleost fishes from coral reefs off eastern Australia and New Caledonia. *Parasite*, 21, 19 <https://doi.org/10.1051/parasite/2014060>
7. Blakeslee, A.M.H., Fowler, A.E. & Keogh, C.L. (2013). Chapter Two - Marine invasion and parasite escape: updates and new perspectives. In: Michael Lesser (ed), *Advances in Marine Biology*, Academic Press, 66, 87-169. <https://doi.org/10.1016/B978-0-12-408096-6.00002-X>
8. Boussellaa, W., Neifar, L., Goedknecht, M.A. & Thieltges, D.W. (2018): Lessepsian migration and parasitism: richness, prevalence and intensity of parasites in the invasive fish *Sphyaena chrysotaenia* compared to its native congener *Sphyaena sphyraena* in Tunisian coastal waters. *PeerJ*, 6, e5558. <https://doi.org/10.7717/peerj.5558>
9. Bush, A.O., Lafferty, K.D., Lotz, J.M. & Shostaka, A.W. (1997). Parasitology meets ecology on its own terms: Margolis et al. Revisited. *Journal of Parasitology*, 83 (4), 575 – 583. <https://doi.org/10.2307/3284227>
10. Castriota, L., Falautano, M., Battaglia, P., Oddo, A. & Andaloro, F. (2014). New biological data on *Fistularia commersonii* in the central Mediterranean Sea. *Cybium - International Journal of Ichthyology*, 38(1), 15-21. <https://doi.org/10.26028/cybium/2014-381-003>
11. CIESM, (2009). Dynamics of Mediterranean deep waters. In: *CIESM Workshop Monographs*, N° 38.

12. Colautti, R.I., Ricciardi, A., Grigorovich, I.A. & MacIsaac, H.J. (2004). Is invasion success explained by the enemy release hypothesis? *Ecology Letters*, 7(8), 721-733 <https://doi.org/10.1111/j.1461-0248.2004.00616.x>
13. Elton, C.S. (1958). *The ecology of invasions by Animals and Plants*. Methuen: London, UK. Springer. <https://doi.org/10.1007/978-1-4899-7214-9>
14. Edelist, D. (2014). New length-weight relationships and L_{max} values of fishes from Southeastern Mediterranean Sea. *Journal of Applied Ichthyology*, 30(3), 521-526. <https://doi.org/10.1111/j.1439-0426.2012.02060.x>
15. Erguden, D., Turan, C. & Gurlek, M. (2009). Weight-length relationships for 20 Lessepsian fish species caught by bottom trawl on the coast of Iskenderun Bay (NE Mediterranean Sea, Turkey). *Journal of Applied Ichthyology*, 25 (1), 133-135. <https://doi.org/10.1111/j.1439-0426.2008.01198.x>
16. Euzet, L. (1994). Order Tetraphyllidea Carus, 1863. In: L. Khalil, A. Jones & R. Bray (eds), *Keys to the cestode parasites of vertebrates*. CAB International, Wallingford, 149-194.
17. Fritzsche, R. A. (1976). A review of the cornetfishes, genus *Fistularia* (Fistulariidae), with a discussion on intrageneric relationships and zoogeography. *Bulletin of Marine Science*, 26(2), 196-204.
18. Gibson, D.I., Jones, A., Bray, R.A. (2002). *Keys to the Trematoda*, Vol. 1. CABI Publishing: Wallingford, UK. [https://doi.org/10.1016/S0035-9203\(03\)90052-6](https://doi.org/10.1016/S0035-9203(03)90052-6)
19. Golani, D. (1998). Impact of Red Sea Fish Migrants through the Suez Canal on the Aquatic Environment of the Eastern Mediterranean. *Bulletin Series Yale School of Forestry and Environmental Studies*, 103, 375-387.
20. Golani, D. (2000). First record of the Bluespotted cornetfish from the Mediterranean. *Journal of Fish Biology*, 56(6), 1545-1547. <https://doi.org/10.1111/j.1095-8649.2000.tb02163.x>
21. Golani, D., Azzurro, E., Corsini-Foka, M., Falautano, M., Andaloro, F. & Bernardi, G. (2007). Genetic bottlenecks and successful biological invasions: the case of a recent Lessepsian migrant. *Biology Letters*, 3, 541-545. <http://doi.org/10.1098/rsbl.2007.0308>
22. Kabata, Z. (2003). Copepods Parasitic on Fishes. In: J.H. Crothers, P.J. Hayward (eds). *Synopsis of British Fauna (New Series)*, Vol. 47 (revised). The Linnean Society & the Estuarine & Coastal Sciences Association, London, 274 pp.
23. Kalogirou, S., Corsini, M., Kondilatos, G. & Wennhage, H. (2007). Diet of the invasive piscivorous fish *Fistularia commersonii* in a recently colonized area of the eastern Mediterranean. *Biological invasions*, 9, 887-896. <https://doi.org/10.1007/s10530-006-9088-3>
24. Kelly, D.W., Paterson, R.A., Townsend, C.R., Poulin, R. & Tompkins, D.M. (2009). Parasite spillback: A neglected concept in invasion ecology? *Ecology*, 90 (8), 2047-2056 <https://doi.org/10.1890/08-1085.1>
25. Koç, H.T. & Erdoğan, Z. (2015). Confirmed occurrence of the long jaw snake eel, *Ophisurus serpens* (Osteichthyes: Ophichthidae) in Edremit Bay (Northern Aegean Sea), Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 15(2) 371-374. https://doi.org/10.4194/1303-2712-v15_2_21
26. Lymbery, A.J., Morine, M., Kanani, H.G., Beatty, S.J. & Morgan, D.L. (2014). Co-invaders: The effects of

- alien parasites on native hosts. *International Journal for Parasitology: Parasites and Wildlife*, 3(2), 171-177. <https://doi.org/10.1016/j.ijppaw.2014.04.002>
27. MacColl, A.D., Chapman, S.M. (2010) Parasites can cause selection against migrants following dispersal between environments. *Functional Ecology*, 24(4): 847-856. <https://doi.org/10.1111/j.1365-2435.2010.01691.x>
 28. Merella, P., Pais, A., Follesa, M.C., Farjallah, S., Mele, S., Piras, M.C. & Garippa, G. (2016). Parasites and Lessepsian migration of *Fistularia commersonii* (Osteichthyes, Fistulariidae): shadows and light on the enemy release hypothesis. *Marine Biology*, 163, 97-97 <https://doi.org/10.1007/s00227-016-2865-3>
 29. Mouine-Oueslati, N., Chater, I., Romdhani, A., Ktari, M.H. & Francour, P. (2017). First biological data on the well established lessepsian migrant *Fistularia commersonii* (Fistulariidae) in the Gulf of Tunis (Central Mediterranean). *Russian Journal of Marine Biology*, 43(69), 503-506. <https://doi.org/10.1134/S1063074017060062>
 30. Muller Ralph, Keys to the trematoda: Volume 1. D. I. Gibson, A. Jones & R. A. Bray (editors). Wallingford: CABI Publishing, 2002. In: Transactions of the Royal Society of Tropical Medicine and Hygiene, 97(1), 2003, 124 p. [https://doi.org/10.1016/S0035-9203\(03\)90052-6](https://doi.org/10.1016/S0035-9203(03)90052-6)
 31. Pais, A., Chessa, L.A., Serra, S., Ruiú, A., Meloni, G. & Donno, Y. (2007). The impact of commercial and recreational harvesting for *Paracentrotus lividus* on shallow rocky reef sea urchin communities in North-western Sardinia, Italy. *Estuarine, Coastal and Shelf Science*. 73 (3-4), 589-597. <https://doi.org/10.1016/j.ecss.2007.02.011>
 32. Petter, A.J., and Maillard, C. (1988). Larves d'Ascarides parasites de poissons en Méditerranée occidentale. *Bulletin du Muséum national d'histoire naturelle. Section A - Zoologie, biologie et écologie animales*, 10(4), 347-369.
 33. Pérez-del-Olmo, A., Kostadinova, A. & Gibson, D.I. (2016). The Mediterranean: high discovery rates for a well-studied trematode fauna. *Systematic Parasitology*, 93, 249-256. <https://doi.org/10.1007/s11230-016-9626-z>
 34. Pipitone, C., D'Anna, G., Coppola, M., Di Stefano, G. & Badalamenti, F. (2004). First record of the Lessepsian fish *Fistularia commersonii* in the western Mediterranean. *Biologia Marina Mediterranea*, 11(3): 327.
 35. Por, F.D. (1978). *Lessepsian migration: the influx of Red Sea biota into the Mediterranean by way of the Suez Canal*. Ecological studies. Springer-Verlag.
 36. Rigby, M.C., Lo, C.M., Cribb, T.H., Euzet, L., Faliex, E., Galzin, R., Holmes, J.C. & Morand, S. (1999). Checklist of the parasites of coral reef fishes from French Polynesia, with considerations on their potential role in these fish communities. *Cybium*, 23 (3), 273-284. <https://doi.org/10.26028/cybium/1999-233-004>
 37. Sanchez-Tocino, L., Hidalgo Puertas, F. & Pontes, M. (2007). Primera cita de *Fistularia commersonii* Ruppell, 1838 (Osteichthyes: Fistulariidae) en aguas mediterráneas de la Península Ibérica. *Zoologica Baetica*, 18, 79-84.
 38. Streftaris, N., Zenetos, A. (2006). Alien marine species in the Mediterranean-the 100 'Worst Invasives' and their impact. *Mediterr. Marine Science*, 7 (1): 87-118.

39. Taskavak, E. & Bilecenoglu, M. (2001). Length-weight relationships for 18 Lessepsian (Red Sea) immigrant fish species from the eastern Mediterranean coasts of Turkey. *Journal of the Marine Biological Association*, 81(5), 895-896.
<https://doi.org/10.1017/S0025315401004805>
40. Zenetos, A., Tsiamis, K., Galanidi, M., Carvalho, N., Bartilotti, C., Canning-Clode, J., Castriota, L., Chainho, P., Comas-González, R., Costa, A.C. (...) (2022). Status and Trends in the Rate of Introduction of Marine Non-Indigenous Species in European Seas. *Diversity*, 14(12), 1077.
<https://doi.org/10.3390/d14121077>

