

SPECIFIC DIVERSITY OF THE BENTHIC MACROFAUNA WITHIN THE WESTERN COAST OF TUNIS BAY AND THE DJERBA ISLAND COAST (SOUTH-WESTERN MEDITERRANEAN)

Ahmed AFLI¹, K. IBIN CHAABANE², R. CHAKROUN³, Ch. JABEUR⁴
and A. A. RAMOS-ESPLÁ⁵

⁽¹⁾ Institut National des Sciences et Technologies de la Mer - 2025 Salammbô – Tunisia

⁽²⁾ Université de Lille 1 – France

⁽³⁾ Faculté des Sciences de Bizerte – Tunisia

⁽⁴⁾ Institut Supérieur de Biotechnologie de Monastir – Tunisia

⁽⁵⁾ University of Alicante, Department of Marine Sciences and Applied Biology – Spain

Corresponding author⁽¹⁾ : Fax: (216) 71 732 622; Email: afl.ahmed@instm.rnrt.tn

التنوع البيولوجي لللافقريات القاعية في الساحل الغربي لخليج تونس وساحل جزيرة جربة (رب البحر الأبيض
(: في هذه الدراسة، تم اقتناء عينات من 21 في سواحل جزيرة جربة و 10 في الساحل الغربي لخليج
. ثم تم بعد ذلك غربلة العينات عبر فتحات مربعة الشكل طول ضلعها 1 ملم، وتم التعرف على اللافقريات وتعدادها.
وفي غياب مؤشر واحد فعال ومناسب لتقييم التنوع البيولوجي، فقد تم اختيار عدة مؤشرات تقليدية. أظهرت النتائج أن
مجتمعات اللافقريات القاعية تختلف في تركيبها اختلافا تاما في المنطقتين، بالرغم من ان عديد الأنواع مشتركة لكنها
تتبادل فيما بينها مراكز الزعامة من منطقة إلى أخرى. لى الفارق الكبير في القيم التي سجلتها المؤشرات
المستخدمة، تم اعتماد مؤشر حسابي معدل يتمثل في احتساب المتوسط
المؤشر المعدل بأن التنوع البيولوجي هو عموما أعلى بكثير في خليج تونس مقارنة بساحل جربة. وتبدو هذه النتا
واقعية بالنظر الى حقيقة مصادر التدهور البيئي ومدى اتساعها في المنطقتين. ففي الواقع، فإن ساحل جربة يخضع في
العقود الأخيرة الى أنشطة حضرية وصناعية متزايدة والى صيد بحري جائر، مما أدى إلى تراجع كبير في الموائل القاعية،
والتي تؤدي بدورها إلى تدهور التنوع البيولوجي.
الكلمات المفتاحية: مؤشرات التنوع البيولوجي، اللافقريات القاعية، مؤشرات التنوع، مؤشرات الهيمنة، السواحل التونسية،

RESUME

Diversité spécifique de la macrofaune benthique dans la côte ouest de la baie de Tunis et la côte de l'île de Djerba (sud-ouest de la Méditerranée) : Dans cette étude, 21 stations ont été échantillonnées, 11 dans la côte de l'île de Djerba et 10 dans la côte ouest de la baie de Tunis. Ensuite, les échantillons ont été tamisés sur une maille carrée de 1 mm de côté, et les animaux capturés ont été identifiés et comptés. Faute d'un indice unique efficace et adapté à toutes les situations, plusieurs indices traditionnels de biodiversité ont été choisis, calculés et discutés. Les résultats montrent que la composition de la communauté est totalement différente dans les deux sites étudiés, et, à chaque site, plusieurs espèces sont communes, mais ils se substituent aux places des leaders d'une station à l'autre. Pour la mesure de la biodiversité et étant donné la grande différence dans les valeurs enregistrées par les indices retenus, une moyenne arithmétique a été calculée à chaque station sur la base des indices les plus cohérents. Elle montre que, globalement, la biodiversité est nettement plus élevée dans la baie de Tunis par rapport à la côte de Djerba. Ces résultats semblent être plus fiables tenant en compte les contraintes réelles environnementales et anthropiques imposées aux deux sites. En effet, la côte de Djerba est l'objet, ces dernières décennies, à de fortes activités urbaines et de pêche industrielle, entraînant une régression significative des habitats benthiques, ce qui peut conduire à la réduction de la biodiversité.

Mots clés: Invertébrés benthiques, indices de biodiversité, indices de richesse, indices de dominance, côtes tunisiennes, indice arithmétique moyen

ABSTRACT

In this study, 21 stations were sampled, 11 in the Djerba island coast and 10 in the western coast of Tunis bay. Then, samples were sifted on a square mesh of 1 mm a side, and the animals collected were identified and counted. Lacking an efficient single index suitable for application in all situations, several traditional nonparametric indices of biodiversity were selected, calculated and discussed. Results show that the community composition is totally different in the two studied sites, and, within each site, several species are common, but

they exchange the leader ranks from one station to another. For biodiversity measurement and since results show varying values depending on indices at several stations, an arithmetic mean index was calculated at each station on the basis of the most similar indices. It shows that, overall, the biodiversity is significantly higher in Tunis bay coast than in Djerba coast. These results are more reliable considering the real environmental / anthropogenic constraints imposed on both sites. Indeed, Djerba coast is subject, these last decades, to strong industrial, urban and fishing activities, causing significant regression of benthic habitats, which can lead to a reduction of the biodiversity.

Keywords : benthic invertebrates, biodiversity indices, richness indices, dominance indices, Tunisian coastal areas, arithmetic mean index

INTRODUCTION

Following the wave of extinction and rarefaction of species during last decades, the concept of "biological diversity" has widely appeared in the scientific literature from the 1970s (Stork 1996, Dubois 2004). Then, the concept of "biodiversity" appeared for the first time in 1985, but has been widely used about 3 years later (Wilson 1988). Several non-parametric indices have been conceived to measure taxonomic biodiversity, their fundamental purpose is to express the data on the number of species and their proportional abundances (Izsák & Papp 2000). Their most important advantages are the ability for direct comparisons between communities that have few or no species in common and the easiness of their application and interpretation (Magurran 2004). These traditional indices fall roughly into three categories; diversity, evenness or dominance indices, according to their mathematical formula weighting more to the species richness or evenness components of community structure (Spatharis & Tsirtsis 2010). Richness indices assume generally a relationship between the number of species and the sample size, dominance indices consider both the number of species and the distribution of the density among them, and evenness indices are simultaneously affected by the total abundance and species richness (Chadwick & Canton 1984, Lamb et al. 2009). Some recent studies have tried to conceive an index able to take into account simultaneously both categories, as weak diversity indices (Ricotta 2002) and the quadratic entropy index (Izsák & Papp 2000). Nevertheless their effectiveness and usefulness need to be confirmed.

Currently, no non-parametric index of biodiversity can be considered ideal and can measure adequately biodiversity in all situations (Clarke & Warwick 1998, 1999, 2001, Snelgrove 1998, Nielsen et al. 2007, Lamb et al. 2009). These traditional indices are yet clearly less efficient to determine the ecological status of marine coastal areas subject to anthropogenic and environmental stresses, and the numerous studies which have tried this have not given real reliable results (Danilov & Ekelund 1999, Foggo et al. 2003, Labruno et al. 2006, 2008, Dauvin et al. 2007). Actually, the biodiversity status of a given

community depends not only on pollution conditions, but also on several other factors (edaphic, hydrodynamic, trophic, etc.) (Afli et al. 2008a, 2009). It is certain that the new biotic indices conceived, for the most of them, in the context of the European Water Framework Directive, specially to assess the ecological status are clearly more appropriate in this examination, because they are based on the sensitivity and the tolerance of species to increasing pollution (Afli et al. 2008b).

The aim of this work is to give a taxonomic knowledge of Tunisian macrofauna, and to study the taxonomic biodiversity of two different Tunisian coastal areas. Indeed, lacking an efficient single index suitable for application in all situations (Lamb et al. 2009), several biodiversity indices on the benthic macrofauna must be used and will be discussed. They can produce reliable results and be more efficient (Taft et al. 2006).

STUDY SITES

The western coast of Tunis bay

Tunis bay covers approximately 350 km² surface, not exceeding 31m depth (Afli et al. 2008a). Its western coast alternates *Posidonia* and *Cymodocea* beds and mud and fine sand substratum. It is subject to industrial and urban development of the northern suburbs of Tunis (Ayari & Afli 2008). For a few decades, significant commercial and fishing activities in ports of Goulette and Radès have been noted, as well as thermal and waste discharges of industrial estates of Radès and of Jebel Jeloud (Diawara et al. 2008). Other pollution sources are also located along the southern and eastern coastline, such as effluents of non-permanent watercourses, food-processing industry, settlements and water-treatment plants which discharge directly into the bay (Ben Charrada & Moussa 1997).

The Djerba island coast

Djerba island covers approximately 500 km² and belongs to the gulf of Gabès. Its coast is subject to the urban waste discharges of Houmet-souk and Ajim settlements and also to the touristic development of Midoun. Whereas, an important industrial, urban and maritime development along the littoral of the gulf of

Gabès (west of Djerba island) has lead to a generalized increase of pollution and impacts on marine systems, producing changes in the structure and functioning of benthic communities (Hamza et al. 2000, Louati et al. 2001, Drira et al. 2008). The gulf of Gabès has great importance for the fishing sector in Tunisia. It contributes about 65% of fishing production and concentrates about 75% of trawlers and almost two thirds of the fishing total fleet (Anonymous 2008).

MATERIAL AND METHODS

In total, 21 stations were sampled for this study, 11 in the Djerba island coast (July 2009) and 10 in the western coast of Tunis bay (May 2008) (figure 1). In the first site where stations are more deep (Table I), three samples were collected at each station by a Van

Veen grab (0.12 m²). In the second site relatively less deep, five samples were collected at each station by diving and using a square metallic quadrat (0.08 m²). In the laboratory, samples were sifted out of the freshwater, on a square mesh of 1 mm a side. Then, the animals collected were preserved with diluted alcohol (70 %) before being identified, for most of them, up to species level.

Obtained data allowed us to calculate at each station the most common biodiversity indices (Table II). A multidimensional analysis was also performed with the Primer software. Specific abundances were transformed using Bray-Curtis similarity of square-root transformed abundance data. Then, a hierarchical cluster analysis (group average mod) was performed. Lacking an ideal biodiversity index, the joint use of several indices to access the taxonomic diversity can give better results, especially if they are of different




Figure 1 : Maps of the study sites showing the location of the sampling stations

Table I : Characteristics of the sampled stations

Sites	Stations	Coordinates (N / E)	Depths (m)	Substrata
Djerba	WJ1	33° 59,018' / 10° 36,929'	20	Sandy muds
	WJ2	33° 56,860' / 10° 38,189'	15,6	Sandy muds
	WJ3	33° 51,018' / 10° 32,737'	22,7	Sandy muds
	WJ4	33° 51,606' / 10° 35,313'	15,6	Sandy muds
	WJ5	33° 52,695' / 10° 38,782'	10,6	Sandy muds
	EJ1	33° 52,501' / 11° 10,219'	32,1	Mud, maerl
	EJ2	33° 49,688' / 11° 13,227'	34,8	Coarse sands
	EJ3	33° 49,802' / 11° 14,406'	39,4	Heterogenous sediments
	EJ4	33° 45,372' / 11° 11,263'	30	Heterogenous sediments
	EJ5	33° 43,762' / 11° 10,469'	25,8	Sandy muds
Tunis	EJ6	33° 42,726' / 11° 08,272'	20,6	Sandy muds
	Goulette	36° 48,833' / 10° 18,641'	< 5	Sandy muds
	Casino	36° 49,041' / 10° 18,871'	< 5	Sandy muds
	Aéroport	36° 49,372' / 10° 19,127'	< 5	Sandy muds
	Khéireddine	36° 49,703' / 10° 19,359'	< 5	Sandy muds
	Kram	36° 50,035' / 10° 19,555'	< 5	Sandy muds
	Salammbô	36° 50,280' / 10° 19,869'	< 5	Sandy muds
	Dermech	36° 50,690' / 10° 19,924'	< 5	Sandy muds
	Dermech Nord	36° 50,964' / 10° 20,203'	< 5	Sandy muds
	Présidence	36° 51,303' / 10° 20,626'	< 5	Sandy muds
	Sidi Boussaïd	36° 52,047' / 10° 21,497'	< 5	Sandy muds

types (richness, abundance, evenness) (Taft et al. 2006, Nielsen et al. 2007, Lamb et al. 2009). Thus, only the indices that show between them a minimum of similarity (grouped together by multidimensional analyses) were used to calculate a single arithmetic mean index (Buckland et al. 2005). Each index has been reduced, divided by the sum of its values at all stations. Then, an average has been established at each station on the basis of these weighted indices. This arithmetic mean index has been, thereafter, calibrated, and three equal intervals were designed corresponding respectively to Low Biodiversity, Moderate Biodiversity and High Biodiversity. This subdivision of the biodiversity into 3 levels may appear subjective, this is because until now there is no ideal reference to which we can refer (Buckland et al. 2005).

RESULTS

The counting of the samples showed that no organism has been found at stations WJ2, WJ4, EJ2 and EJ4. Obviously, these stations were not considered in the analysis. The list of species collected from the sites (Table III) shows that the western coast of Tunis bay is relatively more rich in species (74 species). Gastropods, as *Cerithium scabridum*, *Chrysallida juliae*, *Rissoa spp.*, *Bittium reticulatum* and *Euspira pulchella* represent around 58% of the total number of collected species, followed by polychaetes (30%) as *Phylo foetida*, *Melinna palmata*, *Paraone sp.* and *Glycera spp.* In Djerba island coast, only 13 species were collected. They are distributed into five taxonomic groups (6 polychaetes, 2 bivalves, 2

echnoderms and 2 Crustaceans)., Figure 2 shows that the abundance is relatively

Table II : Summary of characteristics of the indices used to qualify the biodiversity status using benthic communities. S: total number of species; N: total number of individuals; n_i : number of individuals of the species i.

Indices of biodiversity	Algorithms	References
Specific richness	S : number of species	
Margalef's index	$D_{Mg} = \frac{S-1}{Ln(N)}$	Margalef (1958)
Menhinick's index	$D_{Mn} = \frac{S}{\sqrt{N}}$	Menhinick (1964)
Simpson's index	$I' = \sum_{i=1}^S \frac{n_i \times (n_i - 1)}{N \times (N - 1)}$	Simpson (1949)
Shannon-Wiener's index	$H' = -\sum_{i=1}^S \left(\frac{n_i}{N} \times \log_2 \left(\frac{n_i}{N} \right) \right)$	Shannon and Weaver (1963)
Brillouin's index	$HB = \frac{Ln(N!) - \sum_{i=1}^S Ln(n_i!)}{N}$	Brillouin (1962)
McIntosh's index	$U = \sqrt{\sum_{i=1}^S n_i^2}$	McIntosh (1967)
McIntosh's index	$D = \frac{N - U}{N - \sqrt{N}}$	McIntosh (1967)
McIntosh's evenness	$E = \frac{N - U}{N - \frac{N}{\sqrt{S}}}$	McIntosh (1967)
Berger-Parker's index	$d = \frac{n_{max}}{N}$	Berger and Parker (1970)
Pielou's evenness	$J' = \frac{H'}{\log_2(S)}$	Pielou (1966)
Fisher's alpha (α)	$S = a \times Ln \left(1 + \frac{N}{a} \right)$	Fisher et al. (1943)




Figure 2 : Spatial variability of the macrofauna abundance

Table III : List of collected species

Espèces	Djerba	Tunis	Espèces	Djerba	Tunis
<i>Abra alba</i> (W. Wood, 1802)	+		<i>Lyonsia norwegica</i> (Gmelin, 1791)		+
<i>Alvania lineata</i> Risso, 1826		+	<i>Maldane glebifex</i> Grube, 1860		+
<i>Ampelisca</i> sp.	+		<i>Marphysa</i> sp.	+	
<i>Amyclina</i> sp.		+	<i>Melanella</i> sp.		+
<i>Antalis vulgaris</i> (da Costa, 1778)		+	<i>Melarhappe neritoides</i> (Linnaeus, 1758)		+
<i>Aricia foetida imitans</i> Eisig, 1914		+	<i>Melinna palmata</i> Grube, 1870		+
<i>Aricidea (Acmira) cerrutii</i> Laubier, 1966		+	<i>Mitrella minor</i> (Scacchi, 1836)		+
<i>Astropecten</i> sp.	+		<i>Nassarius corniculum</i> (Olivi, 1792)		+
<i>Balanus</i> sp.	+		<i>Nassarius cuvierii</i> (Payraudeau, 1826)		+
<i>Barleeia</i> sp.		+	<i>Nassarius mutabilis</i> (Linnaeus, 1758)		+
<i>Bittium reticulatum</i> (da Costa, 1778)		+	<i>Naticarius</i> sp.		+
<i>Bittium</i> sp.		+	<i>Nephtys</i> sp.	+	
<i>Bolinus brandaris</i> (Linnaeus, 1758)		+	<i>Nereis falsa</i> Quatrefages, 1866		+
Bulimulidae		+	<i>Nereis</i> sp.		+
<i>Calliostoma zizyphinum</i> (Linnaeus, 1758)		+	<i>Notomastus latericeus</i> Sars, 1851		+
<i>Cerithium scabridum</i> Philippi, 1848		+	<i>Nucula nucleus</i> (Linnaeus, 1758)		+
<i>Cerithium</i> sp.		+	<i>Nucula sulcata</i> Bronn, 1831		+
<i>Cerithium vulgatum</i> Bruguiere, 1792		+	<i>Nucula turgida</i> Gould, 1846		+
<i>Chaetozone setoza</i> Malmgren, 1867		+	<i>Nuculana commutata</i> (Philippi, 1844)		+
<i>Chrysallida juliae</i> (de Folin, 1872)		+	<i>Ocenebrina aciculata</i> (Lamarck, 1822)		+
<i>Cirratulus cirratus</i> (O. F. Müller, 1776)		+	<i>Ophiura</i> sp.	+	
<i>Cirriformia tentaculata</i> (Montagu, 1808)		+	<i>Orbinia bioreti</i> (Fauvel, 1919)		+
<i>Columbella rustica</i> (Linnaeus, 1758)		+	Oweniidae	+	
<i>Conus mediterraneus</i> Hwass in Bruguière, 1792		+	<i>Paraonis</i> sp.		+
<i>Corbula gibba</i> (Olivi, 1792)	+		<i>Phyllodoce</i> sp.		+

<i>Ctenocardia</i> sp.		+	<i>Phylo norvegicus</i> (M. Sars in G.O. Sars, 1872)	+
<i>Cucumaria</i> sp.	+		<i>Pisania striata</i> (Gmelin, 1791)	+
<i>Cyclope neritea</i> (Linnaeus, 1758)		+	<i>Prionospio steenstrupi</i> Malmgren, 1867	+
<i>Diaphana minuta</i> T. Brown, 1827		+	<i>Pygospio</i> sp.	+
<i>Euclymene oerstedii</i> (Claparède, 1863)		+	<i>Rissoa paradoxa</i> (Monterosato, 1884)	+
<i>Eulima</i> sp.		+	<i>Rissoa</i> sp.	+
<i>Euspira pulchella</i> (Risso, 1826)		+	<i>Rissoa ventricosa</i> Desmarest, 1814	+
<i>Gastrana fragilis</i> (Linnaeus, 1758)			<i>Ruditapes philippinarum</i> (Adams & Reeve, 1850)	+
<i>Gibberula</i> sp.		+	<i>Sabella pavonina</i> Savigny, 1822	+
<i>Gibbula varia</i> (Linnaeus, 1758)		+	<i>Sabella</i> sp.	+
<i>Glycera convoluta</i> Keferstein, 1862		+	Serpulidae	+
<i>Glycera</i> sp.		+	<i>Smaragdia viridis</i> (Linnaeus, 1758)	+
<i>Glycera unicornis</i> Savigny in Lamarck, 1818		+	<i>Tellina compressa</i> Brocchi, 1814	+
<i>Gyroscala</i> De Boury, 1887		+	Terebellidae	+
<i>Hexaplex (Trunculariopsis) trunculus</i> (Linnaeus, 1758)		+	<i>Tricolia</i> sp.	+
<i>Homalopoma sanguineum</i> (Linnaeus, 1758)		+	<i>Trophonopsis muricata</i> (Montagu, 1803)	+
<i>Kurtiella bidentata</i> (Montagu, 1803)		+	<i>Typhloscolex muelleri</i> Busch, 1851	+
<i>Lumbrineriopsis paradoxa</i> (Saint-Joseph, 1888)		+	<i>Venerupis aurea</i> (Gmelin, 1791)	+
<i>Lumbrineris latreilli</i> Audouin & Milne Edwards, 1834		+		

high at WJ3 (2883 ind./m²) and Dermech Nord (2463 ind./m²), and very low at WJ1, WJ5, EJ1, EJ3, EJ5, EJ6 and Dermech. For the other stations, it varies from 438 ind./m² (Casino) to 1413 ind./m² (Sidi Boussaïd). At first glance, registered values of the biodiversity indices (figure 3) show approximately the same spatial trend, except McIntosh's index U which seems to be very different. Overall, they show low values at stations WJ1, WJ3, EJ5, EJ6, Salammbô and Dermech Nord, high values at stations WJ5, EJ1, EJ3, Aéroport, Khéireddine and Présidence and varying values depending on indices at the other stations. The hierarchical cluster analysis (figure 4) separates, at 57% of similarity, the index U from all

the other indices, which are themselves divided at 73% of similarity into 3 groups. The first group includes H' , D_{Mg} , HB , D_{Mn} and $1/d$, the second includes J' , E , $1-\lambda'$ and D and the third group includes α and S . The calculated values of the arithmetic mean index (AMI) allowed to define 3 equal intervals corresponding to different statuses of biodiversity (figure 5). Thus, 5 stations (WJ1, WJ3, EJ5, EJ6 and Salammbô) were classified in "Low Biodiversity" ($0.000 < AMI < 0.040$), 7 stations (WJ5, EJ1, EJ3, Casino, Kram, Dermech Nord and Sidi Boussaïd) in "Moderate Biodiversity" ($0.040 < AMI < 0.080$) and 5 stations (Goulette




Figure 3 : Calculated values of the biodiversity indices at sampled stations




Figure 4 : Hierarchical clustering (Bray-Curtis similarity) established on calculated values of the biodiversity indices at sampled stations




Figure 5 : The arithmetic mean index calculated on the basis of several biodiversity indices

Aéroport, Khéireddine, Dermech and Présidence) in "High Biodiversity" ($0.080 < AMI < 0.120$). The test of Student applied to the values of AMI in the 2 studied sites shows a significant difference (at $p < 0.005$) in favour of the western coast of Tunis bay. The same test applied separately to each index shows that only the specific richness, Margalef, Shannon-Wiener, Brillouin and Berger-Parker indices indicate significant difference (at $p < 0.005$) between the two studied sites.

DISCUSSION

Although the two study sites are different on a set of characteristics (bathymetry, exposure to anthropogenic activities, hydrological parameters, etc.), used indices show a certain similarity in measuring biodiversity (57% of the Bray-Curtis Similarity). This confirms the robustness of these indices, which should theoretically be applicable to all situations. But in practice, the selection of an index depends on several criteria, such as its

discrimination capacity and sensitivity to sample size, the richness or regularity of the distribution of individuals among different species. In this study for which the difference in sampling effort between the two areas is negligible, the indices used are unanimous in classifying certain stations at biodiversity levels, clearly higher than the average, and some other stations at clearly lower levels. But the problem encountered here concerns stations classified differently, and which represent a good test of robustness for these indices (Afli et al. 2008b). Also, since the interpretation of the absolute value of nonparametric biodiversity indices is subjective, these indices are generally used to compare different communities or the same community over time. Thus, and referring to indices used, the biodiversity seems to be significantly ($p < 0.005$) higher in the eastern coast of Tunis bay than in Djerba coast. On average, in the first site (Tunis bay) an abundance of about 1084 ind./m² are distributed among 74 species, however in the second site (Djerba coast) only about 284 ind./m² are distributed among 13 species.

Nevertheless, among the 12 indices used, only 5 indices (specific richness, Margalef, Shannon-Wiener, Brillouin and Berger-Parker) approve this result. This confirms, once more, that the joint use of a set of indices is usually more efficient than a single index, because generally, the indices have originally been developed to express either the specific richness or proportional abundances of species (Lamb et al. 2009). However, with only data currently available in the literature, we can not give reasonable explanations for these observations. Indeed, the current status of biodiversity is the result of several factors (edaphic, trophic, environmental, human, hydrodynamic, etc.). To estimate the real contribution of each factor, more targeted studies must be undertaken. Nevertheless and referring to works already carried out in the study sites, it seems that Tunis bay despite its exposure to the nuisance of the Tunis city (2250000 inhabitants in 2004), is clearly less subject to environmental / anthropogenic constraints than Djerba coast. Indeed, since a few decades, the gulf of Gabès in general is subject to deep changes, and the 4 azoic stations (WJ2, WJ4, EJ2 and EJ4) confirm these observations. It has been put under anthropogenic pressure due to industrial, urban and fishing activities, causing significant overfishing of demersal resources, the degradation of Seagrass meadows, *Posidonia oceanica*, and the regression of benthic habitats (Zairi & Rouis 1999, Turki et al. 2006). For example, according to Drira et al. (2008), toxic dinoflagellates, mainly *Karenia cf. selliformis*, reach high densities in Djerba coast waters because of the excess reactive nitrogen derived from fertilizer applications, animal wastes and fuel combustion. Moreover, the situation becomes more complicated due to the interaction of various other factors (hydrodynamism, interference in urban areas, shipping, etc.). Actually, all these factors and, certainly others, have induced an important loss of biodiversity in the gulf of Gabès. Chey et al. (1997) and Intachat & Holloway (2000) consider that, besides specific richness, Fisher's alpha is the more efficient biodiversity index to study several communities. In this study, this index is clearly higher only at the station of Dermech where only 8 species and an abundance of 113 ind./m² were registered. Thus, it appears to be not consistent with the arithmetic mean index. Landau et al. (1999) consider that Fisher's alpha and indices of Shannon-Wiener and Simpson, which are the most used measures of diversity in community ecology, are more efficient than the other indices. Nevertheless, the comparison of these indices at, for example, stations of Casino (438 ind./m²) and Salammbô (1275 ind./m²) where 7 species were identified at each of them, show that Fisher's alpha is not consistent with the other indices. Indeed, unlike most used indices, it shows no clear difference between these stations.

Thus, the index alpha can not allow to distinguish between sites that have the same numbers of individuals and of species. It reaches high values for sites dominated by a few species, such as the case of Dermech. This problem is reduced for Shannon-Wiener and Simpson indices, although the Simpson index is sensitive to the abundance of species most dense. As for the dominance index of Berger-Parker, it is considered by May (1975) as the most satisfactory measure of diversity. Nevertheless, this index seems to be different from the other indices, since it expresses the proportional importance of the dominant species. For example, it is the only index that indicates a clear difference between stations of Goulette (14 species and 1225 ind./m²) and Kram (14 species and 1200 ind./m²). According to Magurran (2004), indices of Margalef, McIntosh (U) and specific richness (S) are, in terms of discrimination capacity, the more efficient. However, McIntosh's index (U) seems to be very different from the most other indices, since it takes into account exclusively the abundance. Thus, high values were registered at stations with high abundance, especially at WJ3 and Dermech nord. In general, indices whose the weighting of the specific richness is more important are more useful for detecting differences between sites than indices giving more importance to the dominance / evenness of diversity.

However, Robinson & Tuck (1993) have found that the increase of diversity index with progressive sampling deserves further study to determine after what period diversity stabilizes. In general and according to Magurran (2004), specific richness, Margalef and Menhinick indices are the more sensitive to the sample size, and Simpson and Berger-Parker indices are the less sensitive ones.

In conclusion, the indices of biodiversity used in this study showed different results, among either the two sites or the sampled stations. However, the joint use of these indices gave more reliable results considering the real constraints imposed on both sites. Thus, the arithmetic mean index shows that, overall, the biodiversity is significantly higher in Tunis bay than in Djerba coast.

Acknowledgements

This work was undertaken within the framework of the research activities of the laboratory of "Biodiversité et Biotechnologies Marines" of the INSTM. Many thanks to those who contributed to the field and laboratory components of this study.

BIBLIOGRAPHY

Afli A, Ayari R, Brahim M 2008a. Trophic organization of the macro-zoobenthos assemblages within coastal areas subjected to

- anthropogenic activities. *Journal of the Marine Biological Association of the U.K.*, 88 (4), 663-674.
- Afli A, Ayari R, Zaabi S 2008b. Ecological quality of some Tunisian coast and lagoon locations, by using benthic community parameters and biotic indices. *Estuarine, Coastal and Shelf Science*, 80, 269-280.
- Afli A, Chakroun R, Ayari R, Aissa P 2009. Seasonal and spatial variability of the community and trophic structure of the benthic macrofauna within Tunisian lagoonal and marine coastal areas (south-western Mediterranean). *Journal of Coastal Research*, 25 (6), 1198-1209.
- Anonymous 2008. *Annuaire des statistiques des pêches en Tunisie*. Commissariat Général aux Pêches - Ministère de l'agriculture, Tunisia.
- Ayari R, Afli A 2008. Functional groups to establish the ecological quality of soft benthic fauna within Tunis bay (western Mediterranean). *Vie et Milieu*, 58 (1), 67-75.
- Ben Charrada R, Moussa M 1997. Modélisation hydrodynamique et écologique des eaux côtières du golfe de Tunis. *La Houille Blanche*, 6, 66-78.
- Berger WH, Parker FL 1970. Diversity of planktonic foraminifera in deep-sea sediments. *Science*, 168, 1345-1347.
- Brillouin L 1962. *Science and information theory*. 2nd Ed. Academic Press, New York, USA : 347 pp.
- Buckland ST, Magurran AE, Green RE, Fewster RM 2005. Monitoring change in biodiversity through composite indices. *Philosophical Transactions of the Royal Society B*, 360, 243-254.
- Chadwick JW, Canton SP 1984. Inadequacy of diversity indices in discerning metal mine drainage effects on a stream invertebrate community. *Water, Air and Soil Pollution*, 22, 217-223.
- Chey VK, Holloway JD, Speight MR 1997. Diversity of moths in forest plantations and natural forests in Sabah. *Bulletin of Entomological Research*, 87, 371- 385.
- Clarke KR, Warwick RM 1998. A taxonomic distinctness index and its statistical properties. *Journal of Applied Ecology* 25: 523-531.
- Clarke KR, Warwick RM 1999. The taxonomic distinctness measure of biodiversity: weighting of steps lengths between hierarchical levels. *Marine Ecology Progress Series* 184: 21-29.
- Clarke KR, Warwick RM 2001. A further biodiversity index applicable to species lists: variation in taxonomic distinctness. *Marine Ecology Progress Series* 216. 265-278.
- Danilov R, Ekelund NGA 1999. The efficiency of seven diversity and one similarity indices based on phytoplankton data for assessing the level of eutrophication in lakes in central Sweden. *The Science of the Total Environment*, 234, 15-23.
- Dauvin JC, Ruellet T, Desroy N, Janson AL 2007. The ecological quality status of the Bay of Seine and the Seine estuary: Use of biotic indices. *Marine Pollution Bulletin* 55: 241-257.
- Diawara M, Tlig-Zouari S, Rabaoui L, Ben Hassine OK 2008. Impact of management on the diversity of macrobenthic communities in Tunis north lagoon: systematics. *Cahiers de Biologie Marine*, 49, 1-16.
- Drira Z, Hamza A, Bel Hassen M, Ayadi H, Bouain A, Aleya L 2008. Dynamics of dinoflagellates and environmental factors during the summer in the Gulf of Gabès (Tunisia, Eastern Mediterranean Sea). *Scientia Marina*, 72(1), 59 -71.
- Dubois PJ 2004. *Vers l'ultime extinction? La biodiversité en danger*. Éditions La Martinière, Paris, 191 pp.
- Fisher RA, Corbet AS, Williams CB 1943. The relation between the number of species and the number of individuals in a random sample of an animal population. *Journal of Animal Ecology*, 12, 42-58.
- Foggo A, Rundle SD, Bilton DT 2003. The net result: evaluating species richness extrapolation techniques for littoral pond invertebrates. *Freshwater Biology* 48: 1756-1764.
- Hamza A, Bouain A, El Abed A 2000. Observations sur la floraison et la fructification de la phanérogame marine *Posidonia oceanica* (Linnaeus) Delile sur les côtes du golfe de Gabès (Tunisie). *Mésogée*, 50, 93-99.
- Intachat J, Holloway JD 2000. Is there stratification in diversity or preferred flight height of geometroid moths in Malaysian lowland tropical forest?. *Biodiversity and Conservation*, 9 (10), 1417-1439.
- Izsák J, Papp L 2000. A link between ecological diversity indices and measures of biodiversity. *Ecological Modelling*, 130 (1-3), 151-156.
- Labruno C, Amouroux JM, Sardá R, Dutrieux E, Thorin S, Rosenberg R, Grémare A 2006. Characterization of the ecological quality of the coastal Gulf of Lions (NW Mediterranean). A comparative approach based on three biotic indices. *Marine Pollution Bulletin* 52: 34-47.
- Labruno C, Grémare A, Amouroux JM, Sardá R, Gil J, Taboada S 2008. Structure and diversity of shallow soft-bottom benthic macrofauna in the Gulf of Lions (NW Mediterranean). *Helgolander Marine Research* 62: 201-214.
- Lamb EG, Bayne E, Holloway G, Schieck J, Boutin S, Herbers J, Haughland DL 2009. Indices for monitoring biodiversity change: Are some

- more effective than others?. *Ecological Indicators*, 9, 432-444.
- Landau D, Prowell D, Carlton CE 1999. Intensive versus long-term sampling to assess lepidopteran diversity in a southern mixed mesophytic forest. *Annals of the Entomological Society of America*, 92, 435-441.
- Louati A, Elleuch B, Kallel M, Saliot A, Dagaut J, Oudot J 2001. Hydrocarbon contamination of coastal sediments from the Sfax area (Tunisia), Mediterranean Sea. *Marine Pollution Bulletin*, 42, 445-452.
- Magurran AE 2004. *Measuring Biological Diversity*. Blackwell, Oxford, UK: 256 pp.
- Margalef R 1958. Information theory in ecology. *General Systematics*, 3, 36-71.
- May RM 1975. Patterns of species abundance and diversity (pp 81-120). In : Cody M.L. & Diamond J.M. (eds.), *Ecology and evolution of communities*. Harvard University Press, Cambridge.
- McIntosh RP 1967. An index of diversity and the relation of certain concepts to diversity. *Ecology*, 48, 392-404.
- Menhinick EF 1964. A comparison of some species-individuals diversity indices applied to samples of field insects. *Ecology*, 45, 859-861.
- Nielsen SE, Bayne EM, Schieck J, Herbers J, Boutin S 2007. A new method to estimate species and biodiversity intactness using empirically derived reference conditions. *Biological Conservation*, 137, 403-414.
- Pielou EC 1966. Shannon's formula as a measure of specific diversity : its use and measure. *American Naturalist*, 100, 463-465.
- Ricotta C 2002. Bridging the gap between ecological diversity indices and measures of biodiversity with Shannon's entropy: comment to Izsák and Papp. *Ecological Modelling*, 152, 1-3.
- Robinson GS, Tuck KR 1993. Diversity and faunistics of small moths (microlepidoptera) in Bornean rainforest. *Ecological Entomology*, 18, 385-393.
- Shannon CE, Weaver W 1963. *The mathematical theory of communication*. University of Illinois Press, Urbana, Illinois : 117 pp.
- Simpson EH 1949. Measurement of diversity. *Nature (London)*, 163, 688 pp.
- Snelgrove PVR 1998. The biodiversity of macrofaunal organisms in marine sediments. *Biodiversity and Conservation* 7: 1123-1132.
- Spatharis S, Tsirtsis G 2010. Ecological quality scales based on phytoplankton for the implementation of Water Framework Directive in the Eastern Mediterranean. *Ecological Indicators*, 10, 840-847.
- Stork NE 1996. Measuring global biodiversity and its decline. In *Biodiversity II. Understanding and Protecting our biological Resources* (REAKA-KUDLA M.L., WILSON D.E. & WILSON E.O., dir.), Joseph Henry Press (Washington), 41-68.
- Taft JB, Hauser C, Robertson KR 2006. Estimating floristic integrity in tallgrass prairie. *Biological Conservation*, 131, 42-51.
- Turki S, Harzallah A, Sammari C 2006. Occurrence of harmful dinoflagellates in two different Tunisian ecosystems: the lake of Bizerte and the gulf of Gabes. *Cahiers de Biologie Marine*, 47, 253-259.
- Wilson EO 1988. *Biodiversity*. National Academy Press, Washington, D.C., 538 pp.
- Zairi M, Rouis MJ 1999. Impacts environnementaux du stockage du phosphogypse à Sfax (Tunisie). *Bulletin des Laboratoires des Ponts et Chaussées*, 219, 29-40.